

Fig. 1




Fig.2



# Table 1 結果 爪高指数の推移

	年齢/性別	部位	開始時Start	装着後2週間 2weeks	装着後4週間 4weeks	終了後4週間 after 4weeks	自覚症状Subjective symptoms
症例1	62女Female	右right	55.6%	47.4%	34.7%	41.2%	なし non
症例2		左left	62.1%	41.4%	55.4%	46.4%	なし non
症例3	63女Female	右right	41.0%	39.1%	9.2%	18.6%	直後に消失immediately disappear
症例4		左left	26.4%	17.8%	7.7%	13.3%	直後に消失 Immediately disappear
症例5	59女Female	右right	70.6%	50.0%	24.2%	50.0%	2, 3日で消 disappear after 2or3days
症例6	36男Male	右right	33.3%	11.4%	18.9%	35.3%	なしnon
症例7		左left	44.6%	13.2%	14.7%	33.9%	なしnon
症例8	28女Female	左left	37.9%	17.9%	27.1%	25.7%	2, 3日で消失disappear after 2or3days
症例9	47男Male	左left	76.7%	72.7%	71.1%	90.5%	なしnon
症例10	77女Female	右right	118.8%	73.9%	72.7%	77.3%	1週間で消失 disappear after a week
症例11	52男Male	左left	55.6%	39.6%	28.1%	44.5%	なしnon
症例12	77女Female	右right	51.5%	48.6%	46.5%	41.9%	2, 3日で消失disappear after 2or3days
症例13		左left	82.3%	54.8%	54.2%	45.8%	2週間で消失disappear after a week
症例14	65女Female	右right	56.2%	42.5%	40.5%	50.0%	なしnon
症例15	67女Female	右right	49.5%	31.6%	36.3%	55.0%	なしnon
平均ave			57.5%	40.1%	36.1%	44.6%	

Fig.3



Fig.4



Fig.5

